

Anti-Vibrio from Ethyl Acetate Extract of Sponge- Associated Fungus *Trichoderma longibrachiatum*

Sri Sedjati^{1,4*}, Ambariyanto Ambariyanto^{1,2}, Agus Trianto^{1,2}, Endang Supriyantini¹, Ali Ridlo¹, Ervia Yudiati^{1,3}, Teguh Firmansyah²

¹ Marine Science Department, Faculty of Fisheries and Marine Science, Diponegoro University, Indonesia

² Integrated Laboratory, Diponegoro University, Indonesia

³ Tropical Marine Biotechnology Laboratory, Diponegoro University, Indonesia

⁴ Marine Science Techno Park, Indonesia

ABSTRACT

Some of *Vibrio* spp. bacteria are pathogenic to humans and other organisms, including cultured fish or shrimp. This study aimed to determine the activity of ethyl acetate extract of *Trichoderma longibrachiatum* as an anti-vibrio fungus. The test bacteria used were: *Vibrio harveyi*, *Vibrio anguillarum*, *Vibrio vulnificus*, and *Vibrio parahaemolyticus*. The fungus was cultured using Malt Extract Agar (MEA) medium for 9 days at 27°C (static conditions, 24 hours dark, pH 5.6, and salinity 60 ppt). Extracts obtained by maceration using ethyl acetate, then the extract is partitioned using water-methanol 70:30 and ethyl acetate. Each fraction was concentrated to obtain polar-ethyl acetate (PE) and semipolar-ethyl acetate (SPE) extracts. The components of the constituent extract were traced with a Thin Layer Chromatography (TLC) and followed by ultraviolet spectrophotometry. The anti-vibrio activity was determined based on the value of Minimum Inhibitory Concentration (MIC). The results of the study showed that SPE was more potential to be used as anti-vibrio. The strongest activity was able to inhibit the growth of *Vibrio vulnificus* with 256 µg mL⁻¹ MIC value, while the weakest was against *Vibrio parahaemolyticus* with 1.024 µg mL⁻¹ MIC value. In conclusion, SPE has the potential to be developed as an anti-vibrio compound, particularly against *Vibrio vulnificus*.

Keywords: Semipolar-ethyl acetate extract, Minimum Inhibitory Concentration, *Vibrio vulnificus*.

INTRODUCTION

Vibrio is a genus of bacteria found in a variety of freshwater and marine habitats. There are more than 100 species of *Vibrio* spp., and 12 of which cause infection in humans. *Vibrio cholerae* (*V. cholerae*) can cause cholera, a severe diarrheal disease that can be transmitted through contaminated water. Non-cholera *Vibrio* spp., such as *V. parahaemolyticus*, *V. alginolyticus*, and *V. vulnificus*, may

cause vibriosis, an infection with various clinical expressions, and the mildest of which is gastroenteritis. Vibriosis infection is usually caused by exposure to seawater or by consuming contaminated raw or undercooked seafood. There is an exception, *V. vulnificus* which is an opportunistic pathogen causing wound infection which can rapidly lead to septicemia (1, 2). Infections caused by *V. vulnificus* are generally fatal. Therefore, accurate diagnosis and direct treatment are very important since the infection may cause death to the sufferer (3). antibiotics are widely used to treat vibriosis. The commonly used antibiotics include doxycycline, quinolone, cephalosporin, ciprofloxacin, tigecycline, and

* Corresponding author: Sri Sedjati

sedjati69@gmail.com

Received on 15/1/2021 and Accepted for Publication on 25/8/2021.

ceftazidime (2, 3). *Vibrio* spp. are gram-negative bacteria that causes vibriosis in humans and animals. Moreover, fishery products and post-harvest products are common intermediaries for the transfer of bacteria from their original habitat to their new hosts (5).

The fisheries sector is one of the most fundamental fields for a country's food security. On the other hand, fisheries systems also play a major role in spreading vibriosis to humans. Vibriosis is a bacterial disease reported in Indonesian marine fish culture since the 1990s. The disease is reported mostly found in grouper (*Epinephelus* sp.) and shrimp (*Penaeus monodon*, *Litopenaeus vannamei*) culture, although infection also occurs in snapper (*Lates calcarifer*) and abalone (*Haliotis squamata*) culture. The agent causing vibriosis in marine fish in Indonesia involves several vibrio species, including *V. harveyi*, *V. anguillarum*, *V. alginoluticus*, *V. parahaemolyticus* (4). Some efforts are done to control vibriosis in fish farming activities still rely on the use of drugs or antibiotics given through oral and immersion. However, feed residues will cause pollution in the waters due to antibiotic contamination. Nowadays, several *Vibrio* species are even resistant to certain types of antibiotics. Based on the research (5) conducted on the North Coast of Java found that *V. parahaemolyticus* contaminating vanname shrimp is resistant to erythromycin (90%), amoxicillin-clavulanic acid (83.33%), and nitrofurantoin (58.33%).

Vibrio spp. live in marine habitats along with other bacterial species and microorganisms, like fungi. Competition among the inhabitants of an ecosystem will occur to compete for space and nutrition. Consequently, there is a possibility that there are any antagonistic species, both within among themselves and towards other species. Some research resulted in the fact that several marine-derived fungi species synthesize secondary metabolites of anti-vibrio as their chemical weapons to compete and to avoid predation. Several anti-vibrio metabolites found are such as ethyl acetate extract from sponge-associated fungus *Trichoderma asperellum* (6); prenylxanthone and aspergixanthones from the marine-derived fungus

Aspergillus sp. cultured with shaken Czapek-Dox media (7); indole-diterpenoids and steroids isolated from *Penicillium janthinellum* (8), secondary metabolites of the mangrove-associated fungus *Aspergillus* sp. (9), and secondary metabolites of marine invertebrates associated fungi *A. flavus*, *A. oryzae*, *A. aculeatus*, *Talaromyces minioluteus*, *Hypocrea jecorina*, *Gliomastix murorum*, *Myrothecium inundatum*, and *Curvularia avinis* cultured with Poor Marine Agar (PMA) (10).

The use of antibiotics for a certain period can cause some problem related to pathogenic bacteria resistance towards these antibiotics whether in fish or shrimp body and in addition to the residue which will pollute the environment. Besides, this will also harm humans' health by consuming contaminated sea products. Replacing antibiotics with natural compounds that have antibacterial activity against *Vibrio* spp. offers a new environmentally friendly solution. The potential of sponge-associated fungus *T. longibrachiatum* as a producer of antibacterial compounds has been examined. According to (11), the study results showed that ethyl acetate extract was able to inhibit the growth of several types of both gram-positive and negative bacteria. *Vibrio* is categorized as a member of the gram-negative bacteria group. Accordingly, it is presumed that the extract is also able to inhibit its growth. This study aimed to determine the antibacterial activity of the ethyl acetate extract of sponge-associated fungus *T. longibrachiatum* as anti-vibrio.

MATERIAL AND METHODS

Fungus Isolate

The samples of isolate used in this study were sponge-associated fungi coded TE-PF-03.1. The sponges were collected from the waters of Falajava Beach, Ternate Island, North Maluku, Indonesia (coordinates 00°47'09.12" N; 127° 23'21.76" E) at 3-30 m depth. The fungus has been investigated and identified molecularly using Internal Transcribed Spacer (ITS) rDNA sequence as *T. longibrachiatum* and has been morphologically confirmed (11).

Fungus Culture

Regeneration of fungus collection was carried out before the culture process (i.e. subculture for 7 days). Isolate of *T. longibrachiatum* TE-PF.03.1 was cultured using Malt Extract Agar (MEA Merck) media in 20 Petri dishes. Preparation of MEA was carried out using sterile seawater. According to (11), the final condition of the media showed that the salinity was 60 ppt and pH 5.6. The culture was carried out for 9 days in the environmental condition with 24 hours dark, static, and 27°C temperature.

Secondary Metabolites Extraction

Following 9 day-culture, the media and the micelle were cut into small pieces and macerated with ethyl acetate (1:1v/v). Ethyl acetate extract was obtained after filtration and evaporation processes using a rotary evaporator (40°C) under low pressure. Some part of the ethyl acetate extract was separated for the initial test of its potential as anti-vibrio, while the remaining extract was partitioned using a separatory funnel with water-methanol 70:30 and ethyl acetate (1:1v/v) solvents. Each fraction of polar-ethyl acetate (PE) and semipolar-ethyl acetate (SPE) extract was concentrated and ready to antibacterial test as anti-vibrio (11).

Profiling Secondary Metabolites

The constituents compound of PE and SPE extracts were predicted using the Thin Layer Chromatography (TLC) method (12). In this process, after the extract being developed using a certain mobile phase, it will produce spots on the TLC plate and are identified using R_f (Retention factor) value. The spots on the TLC plate were visualized with ultraviolet (UV) light 365 nm, 2% vanillin-H₂SO₄, 0.25% ninhydrin in acetone, and 1% ferric (III) chloride in methanol (12, 13). Furthermore, the TLC plate was heated at 110°C for 2-3 minutes. Besides, the visualization was done by UV light (λ200-400 nm) which was equipped with an absorption profile using a UV-Vis spectrophotometer.

Antibacterial Activity Test

Vibrio bacteria used in this study were *V. harveyi*, *V. anguillarum*, *V. vulvificus*, and *V. parahaemolyticus*.

The subculture of tested bacteria was carried out in Mueller-Hinton Broth (MHB; Oxoid) and incubated for 24 hours at 37°C. The initial antibacterial bioassay was done by determining the inhibition zone diameter using a disc diffusion assay (14). *Vibrio* bacteria were inoculated in Mueller-Hinton Agar (MHA; Oxoid) media with 0.5 McFarland (1.5x10⁸CFU mL⁻¹) initial density using the swab method. Moreover, the extract preparation was started by making a solution with 50 and 25 mg mL⁻¹ concentration in DMSO. Then, 10 μL of extract solution was dropped onto the surface of sterile disc paper (Oxoid, 6 mm diameter), and it resulted in 500 and 250 μg disc⁻¹ final concentration. The inhibition zone was measured after 24 hours incubation period at 37°C.

Minimum Inhibitory Concentration Test

Minimum Inhibitory Concentration (MIC) test was carried out on fractions PE and SPE of *T. longibrachiatum* ethyl acetate extract. Broth Dilution method was used to determine MIC value referring to the method with resazurin microtiter assay (REMA) (15, 16). The first, 100 μL of the extract solution in DMSO solvent with the highest concentration was filled in the first well in certain rows. Then, the next well was filled with 50 μL of sterile MHB. Moreover, the test material as much as 50 μL was transferred from the first well to the next well to achieve 10 series of dilution (2,048-4 μg mL⁻¹). Meanwhile, 30 μL of resazurin solution (Sigma-Aldrich, 0.02% in distilled water) was added to each well, then 10 μL of *Vibrio* bacterial suspension (1.5 x10⁸ CFU mL⁻¹) were added to each well. Chloramphenicol was used as a positive control (with series concentrations from 64-0.125 μg mL⁻¹). Well the 11th filled DMSO as a negative control and the 12th filled MHB as media control. The microplate was incubated at 37°C for 24 hours. After the incubation period, the well as growth control appeared pink. At last, the MIC value was determined based on the lowest concentration that could inhibit the growth of *Vibrio* spp. and appeared blue.

RESULTS AND DISCUSSION

Based on the results of previous studies, *T. longibrachiatum* cultured using MEA media will reach the peak of its secondary metabolites production on day 7. At the following periods, the production decreased, but the antibacterial activity increased until the 9th day of culture. The ethyl acetate extract metabolite has antibacterial activity through the disc diffusion test (500 µg disc⁻¹) against several gram-positive and negative bacteria, while the methanol extract is inactive (11). Similar results were also found by other researchers stating that the ethyl

acetate metabolite extract from endophytic fungal isolates mostly showed higher antibacterial activity than the methanol extract (17).

T. longibrachiatum sponge-associated fungus in this study was cultured using MEA for 9 days. Besides, the anti-vibrio potential of its ethyl acetate extract on disc diffusion test (500 and 250 µg disc⁻¹) can be seen in Table 1 and it shows a inhibition zone formed. This ethyl acetate extract was able to inhibit the growth of *V. harveyi*, *V. anguillarum*, *V. vulvificus*, and *V. parahaemolyticus*.

Table 1. Anti-vibrio potential of *T. longibrachiatum* ethyl acetate extract by using disc diffusion method

Tested Bacteria	Concentration of extracts (µg disc ⁻¹)	
	500	250
<i>V. harveyi</i>	+	-
<i>V. anguillarum</i>	+	+
<i>V. vulvificus</i>	+	+
<i>V. parahaemolyticus</i>	+	+

Note: += inhibition zone formed, - = no inhibition zone

Further research was done to determine the polarity of the active compounds carried out by partition method using immiscible solvents. Methanol is added water to increase its polarity, so that it can be separated with ethyl acetate. The profile of the compounds in PE and SPE extracts corresponded to the appearance of several spots on the TLC plate as seen in Figure 1. The spots appeared fluorescent blue when being irradiated by UV indicating that the organic compound had double bonds (diene/polyene or conjugated) (18). Compounds that react positively with vanillin reagent would show certain colored spots (varying colors) characterizing the presence of carbonyl functional groups (ketones, aldehydes). These

compounds probably come from phenolic, flavonoids, terpenoids, steroids, fatty acids/essential oils, or high molecular weight alcohol groups. Besides, vanillin reagent is sensitive to the steroid class. If the compound reacts positively with the ninhydrin reagent, the compound contains nitrogen (amines, peptides, or alkaloids). Meanwhile, phenolic group compounds will react positively to ferric (III) chloride reagent (12, 19). The results of the study showed that all spots in ethyl acetate extract were not reactive to ninhydrin and ferric (III) chloride reagents; it is accordingly assumed that they are not from nitrogen or phenolic compounds. Perhaps both extract PE and SPE are terpenoids or steroids.




Figure 1. TLC profile of *T. longibrachiatum* ethyl acetate extract and its visualization using: (a) 0.25% ninhydrin, (b) 1% ferric (III) chloride, (c) UV light 365 nm, (d) 2% vanillin-H₂SO₄ (Note: PE=polar-ethyl acetate, SPE=semipolar-ethyl acetate)




Figure 2. UV spectra (λ 200-400 nm) of *T. longibrachiatum* ethyl acetate extract: (a) PE=polar-ethyl acetate , (b) SPE=semipolar-ethyl acetate)

The spectra of UV light absorption is shown in Figure 2. Extract PE has a pattern with 1 peak, i.e. at λ 267.6 (A=0.93), while SPE has 2 clear peaks, at λ 255.2 nm (A=4.00) and λ 276.8 (A=0.80). There may be other peaks in SPE. This was also reflected in the TLC profile of SPE which was observed in more than 2 spots after visualization with 2% vanillin-H₂SO₄ (as in Figure 1d). Based on the literature study, some terpenoids and steroids possess a carbonyl functional group and also conjugated double bonds. This assumption is strengthened by

observing the UV spectra patterns of each PE and SPE. The result of previous research (20), the UV spectra of terpenoids seems to have an absorption peak at λ 259.57 nm for monoterpenoids (C₁₀) compounds which are assumed to be thymol and at λ 260.47 nm for sesquiterpenoids (C₁₅) which is assumed to be chiloscyphone. The structure of chiloscyphone compounds contains ketone functional groups and conjugated double bonds, whereas thymol only has conjugated bonds. This assumption was strengthened by previous statements (21), that

conjugated double bond which also has a carbonyl functional group will produce chromophores that intensively absorb UV light in λ 230-270 nm range, and weakly at 300-330 nm. Meanwhile, regarding the chemical structure of steroids, 2 double bonds can be distributed between two adjoining rings (heteroannular diene). In this case, the steroid will absorb in the UV region at 220-250 nm. Moreover, it is also possible that 2 ethylene bonds ($H_2C = CH_2$) are in the same ring (homoannular dienes). Consequently, this will shift the absorption peak to 260-285 nm.

Based on the data in the following Table 2, the SPE

obtained from *T. longibrachiatum* has better anti-vibrio activity than the PE. It has strongest potential against *V. vulviniificus* with a $256 \mu g mL^{-1}$ MIC value. There are only 2 species sensitive to chloramphenicol, i.e. *V. harveyi* and *V. vulviniificus*, and 2 others that are categorized in the intermediate group. The anti-vibrio potential of SPE is still weak; this is presumably because the extract is not yet pure. The extract is still in the form of a mixture of several compounds which may not be fully synergistic in supporting its antibacterial properties.

Table 2. MIC values of polar-ethyl acetate (PE) and semipolar-ethyl acetate (SPE) extracts against *Vibrio* spp.

Tested Bacteria	MIC value against <i>Vibrio</i> spp. ($\mu g mL^{-1}$)		
	PE	SPE	Chloramphenicol*
<i>V. harveyi</i>	> 2,048	512	8
<i>V. anguillarum</i>	> 2,048	512	16
<i>V. vulviniificus</i>	1,024	256	2
<i>V. parahaemolyticus</i>	1,024	1,024	16

Note: * Categories on organism as susceptible (22):

MIC>32 = resistant, 16-32 =intermediate, <8 =sensitive

Based on various results of other previous studies, there are many terpenoids isolated from fungi. *Sesquiterpenes*, *meroterpenes*, and *diterpenes* make up the largest proportion of *terpenes*. The genera of *Penicillium*, *Aspergillus*, and *Trichoderma* fungi are terpenoid's dominant producers. The majority of fungi isolated from living material from the sea (animals and plants) produce terpenoids and many exhibit various bioactivities, such as cytotoxicity, anti-inflammatory, enzyme inhibitors, including antibacterial activities (23, 24). Similarly in butanolic extract of terrestrial *Trichoderma* sp. (isolated from forest plants), it is mostly dominated by terpenoid compounds identified as terpenes (limonene) (92.6%). Other constituents represent a small amount proportion consisting of hydrocarbons (2.01%), alcohol (2.4%), ketones (1.78%), and esters (1.03%). Culture activity using Potato Dextrose Agar (PDA) media produces more

terpenoid metabolites than MEA. The growth of *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Micrococcus luteus* was inhibited at $500 \mu g mL^{-1}$ concentration, while the growth of *Escherichia coli* was inhibited at $1 mg L^{-1}$ concentration (25).

The compounds in extract SPE were relatively less polar; as the character of terpenoids or steroids which are composed of isoprene (C5) hydrocarbon and were presumed to have a carbonyl or hydroxyl group. The presumption of antimicrobial properties is according to (26) stating that oxygenated terpenes show strong antibacterial activity, particularly against gram-negative bacteria. The increase in antimicrobial activity is associated with the presence of hydroxyl functional groups (phenolic compounds or alcohols), while the hydrocarbon groups produce relatively low activity. The bacteria which could be inhibited were *S. aureus*, *Bacillus cereus*, *E. coli*, and *Salmonella enterica*.

Some oxygenated terpene compounds such as carvacrol, l-carveol, eugenol, trans-geraniol, and thymol showed higher activity when compared to sulfanilamide. Terpineol showed excellent bactericidal activity against *S. aureus* strains. Meanwhile, carveol, citronellol, and geraniol exerted a rapid bactericidal effect against *E. coli*. The images obtained by scanning electron microscopy (SEM) show that the mechanism which causes the death of the bacterial cell is based on the loss of integrity on cellular membrane function.

The steroid group and its derivatives isolated from several strains of fungi have also been examined and the results of the study showed antimicrobial activities. Steroids are also produced by *Trichoderma sp.*, *Penicillium sp.*, and *Acremonium sp.* with their greatest abundance respectively are ergosterol, ergostatetraenol, ergostapentaene, neoergosterol, and eburicol. Ergosterol is a sterol (alcohol steroid) commonly found in the plasma membrane of fungi. These steroids have antimicrobial activity against several gram-positive and negative bacteria, for instance. *S. aureus*, *Bacillus sp.*, *B. cereus*, *Listeria ivanovii*, *E. coli*, *Citrobacter freundii*, and *Salmonella* spp. However, their antimicrobial activity is weak

against all bacteria with highest inhibition zone for eburicol (24 mm) (27). Similar results were also seen in this study, extract SPE was able to inhibit the growth of *V. harveyi*, *V. anguillarum*, *V. vulviniificus*, and *V. parahaemolyticus*, but with high MIC values, ranging from 256-1,024 $\mu\text{g mL}^{-1}$.

CONCLUSION

Semipolar-ethyl acetate (SPE) extract from sponge-associated fungus *T. longibrachiatum* has stronger antibacterial activity than its polar-ethyl acetate (PE), so it is concluded that SPE has the potential to be developed as an anti-vibrio compound. Besides, the strongest potential of SPE was able to inhibit the growth of *V. vulviniificus* with 256 $\mu\text{g mL}^{-1}$ MIC value, while the weakest was against *V. parahaemolyticus* with 1,024 $\mu\text{g mL}^{-1}$ MIC value.

ACKNOWLEDGEMENT

This scientific paper was written based on a research study supported by the Grant Program of Faculty of Fisheries and Marine Science, Diponegoro University. The funding was intended for research activities from the Fiscal year 2020 with contract number: 026/UN 7.5.10.2/PP/2020.

REFERENCES

- (1) Huehn S., Eichhorn C., Urmersbach S., Breidenbach J., Bechlars S., Bier N., Alter T., Bartelt E., Frank C., Oberheitmann B., Gunzer F., Brennholt N., Böer S., Appel B., Dieckmann R. and Strauch, E. Pathogenic vibrios in environmental, seafood and clinical sources in Germany. *Int J Med Microbiol* 2014; 304(7): 843–50.
- (2) Baker-Austin C., Oliver J.D., Alam M., Ali A., Waldor M.K., Qadri F. and Jaime Martinez-Urtaza J. *Vibrio* spp. infections. *Nat Rev Dis Prim* 2018; 4(1): 8-19.
- (3) Yu W., Shen X., Pan H., Xiao T., Shen P. and Xiao Y. Clinical features and treatment of patients with *Vibrio vulnificus* infection. *Int J Infect Dis* 2017; 59:1–6.
- (4) Istiqomah I., Sukardi S., Murwantoko M. and Isnansetyo A. Review vibriosis management in Indonesian marine fish farming. *E3S Web Conf* 2020; 147: 01001
- (5) Kusmarwati A., Yennie Y. and Indriati N. Antibiotic resistance in *Vibrio parahaemolyticus* from Vannamei shrimp originated from Northern Coast of Java for export market. *J Pascapanen dan Bioteknologi Kelaut dan Perikanan* 2017; 12(2):91–105.
- (6) Sibero M.T., Radjasa O.K., Sabdono A., Trianto A., Triningsih D.W. and Hutagaol I.D. Antibacterial activity of Indonesian sponge associated fungi against clinical pathogenic multidrug resistant bacteria. *J Appl Pharm Sci* 2018; 8(2):088–94.
- (7) Zhu A., Zhang X.W., Zhang M., Li W., Ma Z.Y., Zhu H.J. and Cao F. Aspergixanthonones I–K, new anti-vibrio prenylxanthonones from the marine-derived fungus *Aspergillus* sp. ZA-01. *Mar Drugs* 2018; 6(9):1–8.
- (8) Guo X.C., Xu L.L., Yang R.Y., Yang M.Y., Hu L.D., Zhu

- H.J. and Cao F. Anti-vibrio indole-diterpenoids and C-25 epimeric steroids from the marine-derived fungus *Penicillium janthinellum*. *Front Chem* 2019; 7:1–7.
- (9) Haryani Y., Hilma R., Delfira N., Martalinda T., Puspita F., Friska A., Juwita D. and Ardi F. Anti-vibriosis activity of endophytic fungi associated with *Cerriops tagal* (Perr.) C.B.Rob and *Bruguiera* sp., mangrove plants from Riau Province, Indonesia. *AIP Conf Proc* 2020; 2243(1): 030006.
- (10) Trianto A., Radjasa O.K., Sibero M.T., Sabdon A., Haryanti D., Zilullah W.O.M., Syanindyta A.R, Bahry M.S., Widiananto P.A., Helmi M., Armono H.D., Supriadi S. and Igarashi Y. The effect of culture media on the number and bioactivity of marine invertebrates associated fungi. *Biodiversitas* 2020; 21(1):407–412.
- (11) Sedjati S., Ambariyanto A., Trianto A., Supriyantini E., Ridlo A., Bahry M.S., Wismayanti G., Radjasa O.K. and McCauley E. Antibacterial activities of the extracts of sponge-associated fungus *Trichoderma longibrachiatum* against pathogenic bacteria. *Squalen Bull Mar Fish Postharvest Biotechnol* 2020; 15(2): 81–90.
- (12) Harborne J.B. *Phytochemical Methods; A Guide to Modern Techniques of Plant Analysis*. 1998. 317 p.
- (13) Sen S., Sarkar S., Kundu P. and Laskar S. Separation of amino acids based on Thin-Layer Chromatography by a novel quinazoline based antimicrobial agent. *Am J Anal Chem* 2012; 3(9):669–674.
- (14) Trianto A., Widyaningsih S., Radjasa O.K. and Pribadi R. Symbiotic fungus of marine sponge *Axinella* sp. producing antibacterial agent. *IOP Conf Ser: Earth Environ Sci* 2017; 55: 012005.
- (15) Sowjanya P., Srinivasa B.P. and Lakshmi N.M. Phytochemical analysis and antibacterial efficacy of *Amaranthus tricolor* (L) methanolic leaf extract against clinical isolates of urinary tract pathogens. *African J Microbiol Res* 2015; 9(20):1381–1385.
- (16) Fajarningsih N.D., Munifah I. and Zilda D.S. Evaluation of Antibacterial assays for screening of marine invertebrate extracts. *Squalen Bull Mar Fish Postharvest Biotechnol* 2018; 13(1):1–8.
- (17) Choudhary, M.I., Mohammad, M. Y., Musharraf, S. G. and Atta-ur-Rahman. Epoxidation of ferutin by different fungi and antibacterial activity of its metabolite. *Jordan J. Pharm. Sci.* 2013; 6 (1): 23–29.
- (18) Shadid, K. A., Al-Lahham, S., Jaradat, N., Abu-Nameh, E. S. M. and Qaisi, A. M. Preliminary phytochemical screening, antioxidant and antimicrobial activities of the aqueous, methanol, acetone, and hexane fractions of *Centaurea cyanoides* Wahlenb. *Jordan J. Pharm. Sci.* 2019; 12 (1): 21–32.
- (19) Leylaie S. and Zafari D. Antiproliferative and antimicrobial activities of secondary metabolites and phylogenetic study of endophytic *Trichoderma* species from Vinca plants. *Front Microbiol* 2018; 9:1–16.
- (20) Mohammed A.M. UV-Visible spectrophotometric method and validation of organic compounds. *Eur J Eng Res Sci* 2018; 3(3): 8–11.
- (21) Jork H, Funk W, Fischer W, Wimmer H, Burns DT. *Thin-Layer Chromatography-Reagents and Detection Methods, Physical and Chemical Detection Methods: Fundamentals, Reagents I. Volume 1a*. 1990; 511–512 p.
- (22) Patel M.R., Panchal H.S. and Saluja A.K. Identification of terpenoids and steroidal compounds in *Caryota urens* leaves by column chromatography and various spectroscopic techniques. *World J Pharm Pharm Sci* 2016; 5(5): 1610–1622.
- (23) Dorfman L. Ultraviolet absorption of steroids. *Chem Rev* 1953; 53(1):47–144.
- (24) Dolinsky A.L. Performance Standards for Antimicrobial Susceptibility Testing. *M100* 2017: 27–39.
- (25) Jiang Z, Kempinski C. and Chappell J. Extraction and analysis of terpenes/terpenoids. *Curr Protoc Plant Biol* 2016 ; 1: 345–358.
- (26) Jiang M., Wu Z., Guo H., Liu L. and Chen S. A review of terpenes from marine-derived fungi: 2015–2019. *Mar Drugs* 2020; 18(6): 2015–2019.
- (27) Khethr F.B.H., Ammar S, Saidana D, Daami M., Chriaa J., Liouane K., Mahjoub M.A., Helal A.N. and

- Mighri Z. Chemical composition, antibacterial and antifungal activities of *Trichoderma* sp. growing in Tunisia. *Ann Microbiol* 2008; 58(2): 303–308.
- (28) Guimarães A.C., Meireles L.M., Lemos M.F., Guimarães M.C.C., Endringer D.C., Fronza M. and Scherer R. Antibacterial activity of terpenes and terpenoids present in essential oils. *Molecules* 2019; 24(13): 1–12.
- (29) Dos Santos Dias A.C., Couzinet-Mossion A., Ruiz N., Lakhdar F., Etahiri S., Bertrand S., Ory L., Roussakis C., Pouchus Y.F., Nazih E.H. and Wielgosz-Collin G. Steroids from marine-derived fungi: Evaluation of antiproliferative and antimicrobial activities of eburicol. *Mar Drugs* 2019; 17(6): 372-377.

المضادة للفيبريو من استخراج خلات إيثيل من جمعية الفطر-الإسفنج *Trichoderma longibrachiatum*

سري سيدجاتيو^{1,4,*}، أمبريانتو أمبرانته^{1,2}، أغوس، ترانته^{1,2}، علي ريدلو¹، إندانغ سوبريانتي¹،
إرفيا يودي^{1,3}، تيغوه فيرمانسيه²

¹ قسم العلوم البحرية، كلية المصايد والعلوم البحرية، جامعة ديونيجورو، اندونيسيا

² المختبرات المدمجة، جامعة ديونيجورو، اندونيسيا

³ مختبر علم الأحياء البحرية الاستوائية، جامعة ديونيجورو، اندونيسيا

⁴ مارين ساينس تكنو بارك، جامعة ديونيجورو، اندونيسيا

ملخص

بعض أنواع البكتيريا *Vibrio spp.* هي مسببات الأمراض للبشر والكائنات الحية الأخرى، بما في ذلك الأسماك المستزرعة أو الروبيان. هدفت هذه الدراسة إلى تحديد فعالية مستخلص أسيتات الإيثيل من *Trichoderma longibrachiatum* مثلًا لمضادة لـ *Vibrio spp.* كانت بكتيريا الاختبار المستخدمة هي: *Vibrio harveyi*، *Vibrio parahaemolyticus*، *Vibrio vulvificus*، *Vibrio anguillarum*، تمت زراعة الفطر باستخدام وسط مستخلص الشعير لمدة 9 أيام عند 27 درجة مئوية (ظروف ثابتة، 24 ساعة مظلمة، درجة الحموضة 5.6، و الملوحة 60 لكل ألف). يتم الاستخراج الأولي عن طريق التكسير باستخدام خلات الإيثيل، ثم تتركز. يتم تقسيم المقتطف مع المذيبات الميثانول أكواديس (70:30) وخلات الإيثيل بنسبة 1:1. القطبية الإيثيل. ويتركز كل جزء بحيث يتم الحصول على مقتطفات خلات الإيثيل القطبي ومستخلصات خلات شبه القطبية الإيثيل. يتم تتبع مكونات المركبات المكونة لها باستخدام طريقة الكروماتوغرافيا طبقة رقيقة. نشاط المضادة لـ *Vibrio* يتم تحديده على أساس الحد الأدنى لقيمة التركيز المثبطة باستخدام اختبار ريسازورين. وأظهرت النتائج أن مقتطفات خلات شبه القطبية الإيثيلية هي أكثر قدرة على مكافحة *Vibrio*. أقوى نشاط لها قادر على منع نمو *V. vulvificus* مع الحد الأدنى من قيمة تركيز المثبطة من 256 جزء في المليون، في حين أن أضعف ضد *V. parahaemolyticus* بقيمة 1024 جزء في المليون. المركبات التي يعتقد أنها بمثابة المضادة للفيبريو هي تيربينويدات أو المنشطات. في الختام، يمكن تطوير خلات استخراج الإيثيل شبه القطبي كمركب مضاد للفيبريو، وخاصة ضد *V. vulvificus*.

الكلمات الدالة: استخراج خلات شبه القطبية الإيثيلية، الحد الأدنى من التركيز المثبط، *Vibrio vulvificus*.

* المؤلف المراسل: سري سيدجاتيو

sedjati69@gmail.com

تاريخ استلام البحث 2021/1/15 وتاريخ قبوله للنشر 2021/8/25.